FIG WORKING WEEK 2019

22–26 April, Hanoi, Vietnam

"Geospatial Information for a Smarter Life and Environmental Resilience"

ORGANISED BY FIG

PLATINUM SPONSORS esri THE SCIENCE OF WHERE Trimble

FIG WORKING WEEK 2019
22–26 April, Hanoi, Vietnam
"Geospatial Information for a Smarter Life and Environmental Resilience"

The Potential of Using Satellite Altimetry for Detecting Sea Level Changes in Brunei

Muhammad Hifney Hj Abd Rahman
Surveyor
Brunei Survey Department

ORGANISED BY FIG

PLATINUM SPONSORS esri Trimble

CONTENTS

- Introduction
- Objectives & Outcomes
- Background
- Satellite Altimetry for Sea Level Study
- Methodology
- Results & Analysis
- Further Work
- Conclusion

INTRODUCTION

- The rise of sea level change raised concerns to the earth's population
- Global sea level rise is accelarating incrementally over time in the last 25 years
- Two major factors:
  - Thermal expansion
  - Melting of glaciers and ice sheets

AIMS

- To investigate the potential of using satellite altimetry data for studying sea level changes in Brunei coastal area

OBJECTIVES & OUTCOMES

- Understand the basic and fundamental concepts of satellite altimetry
- Review on previous study
- How altimetry data been used and challenges
- Data plotting using Phyton Programming software from Jason-1 and Jason-2 mission
- Data assessment with recommendations on the reliability of the data

HOW SEA LEVEL CAN BE DETERMINED?

- Tide Gauge and Global Navigation Satellite System (GNSS) – effected by vertical land motion (VLM)

- Satellite Altimetry – satellite-based technology with reference to earth's center (independent)

FACTORS OF SEA LEVEL CHANGES

- Thermal Expansion – rise in sea surface temperature
  - Ocean warmed by 0.009°C to 0.13°C per decade (IPCC, 2014)
- Sea level rise in coastal area due to tides and storm surge
- Vertical Land Motion
- Thermohaline Circulation – water density increased due to temperature and salinity

GLOBAL SEA LEVEL TREND

- Rise of sea level concerned the world's population especially in the coastal areas
- Sea level rise can cause flooding, faster rate of erosion of cliffs and beaches and permanent submersion
- Sea level predicted to keep on rising in the next decade
- Global sea level rise at 1.6mm-1.8mm per year (tide gauge)
- 3.2mm-3.4mm per year since 1992 (satellite altimetry)

GLOBAL SEA LEVEL TREND



25 years of multi-mission sea level trend from altimetry (Source: ESA and CNES/LEGOS (AVISO), 2017)

IMPORTANCE OF SEA LEVEL STUDIES

- Disaster mitigation plan
- Coastal management

IMPORTANCE OF SEA LEVEL STUDIES



Regions that are vulnerable to coastal flooding due to sea level rise (Source: Nicholls and Cazenave, (2010))

IMPORTANCE FOR BRUNEI DARUSSALAM?

- Geographical Location
- Industrial projects on islands and coastal region
- Long-term mitigation plan

PREVIOUS STUDY IN SOUTH CHINA SEA

- World Bank Group published sea level anomaly for South China Sea based on T/P mission indicated a rise between 1992-2008
- Study by Li (2002) found there is a rise at the rate of 10mm/year with warming rate at 0.15° C/ year between 1993-1999
- Sea level fall in 1997-1998 due to El-Nino event (Cheng and Qi, 2007)
- Latest study by Hamid et.al (2016) indicated a rise at 3.85 mm/year (1993-2005) by using multi mission

PRINCIPLES OF SATELLITE ALTIMETRY



PRINCIPLE OF SATELLITE ALTIMETRY

- Measurement of time taken by a radar pulse to travel from satellite to the sea surface and back to the satellite
- Satellite location based on latitude, longitude and satellite altitude coordinates
  - GNSS Satellites
  - DORIS station
  - Satellite Laser Ranging (SLR)

ERRORS AND CORRECTIONS

- Waves pass through the atmosphere can be decelerated by water vapour and inonisation
- Corrections:
  - Range corrections
  - Geophysical corrections – tides and atmospheric pressure

SATELLITE ALTIMETRY MISSION

@NASA JPL(2018)

SATELLITE ALTIMETRY FOR SEA LEVEL

- Since 1992
- Better accuracy when compared to tide gauge - independent



Source: climate.nasa.gov

PROJECT OVERVIEW



DATA SOURCE

- Geophysical Data Record (GDR) – AVISO
- Data contains sensor measurements and full set of geophysical corrections
- Two sets
  - Radar Altimetry Database System (RADS)
  - Sensor Geophysical Data Record (SGDR)
- Downloaded from NASA

RADAR ALTIMETRY DATABASE SYSTEM (RADS)

- Provides simplification for reading, editing and handling
- Users able to access to the up-to-date range and geophysical corrections
- Consists of 1 Hz waveform – 1 point every 6 kilometres

SENSOR GEOPHYSICAL DATA RECORD (SGDR)

- Full accuracy altimeter
- High precision orbit
- Accuracy approximately 1.5 cm
- Contains all relevant corrections for sea surface height calculation
- Include 20Hz waveform – 1 point every 300m

DATA

- Processed data from Jason I and Jason II mission between 2002-2016
- Tracking path near Brunei coastline
- Jason mission tracks every 10 days during its mission
- Approximately 550 tracks


Jason Mission Satellite Tracks over Brunei Coastline
113°0'0"E
114°0'0"E
115°0'0"E
116°0'0"E
4.0.0.6 N
4.0.0.8 N
4.0.0.7 N
4.0.0.9 N
4.0.0.5 N
4.0.0.4 N
4.0.0.3 N
114
038
051
WILAYAH PERSEKUTUAN LABUAN
Bandar Seri Begawan
BRUNEI
BELAIT
MALAYSIA
SARAWAK
N
Legend
Satellite Tracks

DATA PROCESSING

- Processed by Dr Nadim Dayoub – National Oceanography Centre, Southampton
- Sea surface height processed relative to DTU15MSS model then computed using ALES retracker
- DTUMSS15? Latest release model for high resolution mean sea surface
- Time series sea level trend calculated using Robust Regression analysis

DATA PLOTTING

- Time series sea level trend calculated using Robust Regression analysis
- Data in netCDF format
- Python programming – ability to handle and plot netCDF data

RESULTS AND ANALYSIS

- Sea level trends from 2002-2016 from three satellite tracks near Brunei coastline



RESULTS AND ANALYSIS

- Average sea level trend rise at 5.5 mm/year



RESULTS AND ANALYSIS



DISCUSSION AND RECOMMENDATION

- Sea Level Trend
  - The sea level trend in this project does agree with previous study
- Limitations
  - Extensive period can give clear indication of the trend
  - Satellite coverage not exactly at Brunei coastline
  - Errors as it approached the coastline

FUTURE WORK

- Data validation against existing tide gauges – one tide gauge collected data since 1990
- Vertical land motion study – availability of tide gauges and GNSS
- Study of water level for rivers and lakes
- Sentinel-3 mission – extensive satellite coverage

FUTURE WORK – SENTINEL 3 MISSION

- First launched in 2016



WAY FORWARD

- Sea level trend prediction in the next 10 years
- Information from this project – early mitigation plan
- Further investigation of sea surface temperature
- Altimetry data processing expertise

CONCLUSION

- Study able to predict the sea level trend of Brunei Darussalam
- Sea level trend for Brunei Darussalam rising approximately at the rate of 5.5mm/year between 2002-2016
- Satellite altimetry has high potential for sea level study in Brunei Darussalam

THANK YOU!

hifney.rahman@survey.gov.bn